

This leaflet is to tell you about **Salmonella infection**. Although this illness usually doesn't cause any long term problems, it is an unpleasant condition while it lasts.

Now read on...

What is Salmonellosis and how common is it?

Salmonellosis is the term for an infection with a bacterium (germ) called Salmonella.

Salmonellosis is more common in the summer than in the winter and many mild cases are not diagnosed or reported.

How is Salmonellosis caught?

- Salmonella lives in the guts of humans and other animals, including birds, and is usually transmitted to humans by eating contaminated food. The contaminated foods look and smell normal.
- Salmonella is found in foods such as poultry, beef and eggs but all foods including vegetables can become contaminated. Raw foods pose a greater threat than cooked foods as thorough cooking kills Salmonella.
- Food may become contaminated by infected food handlers who do not wash their hands after going to the toilet.
- Salmonella may also be found in the faeces of some pets, especially those with diarrhoea, and people can become infected if they do not wash their hands after contact with these faeces. 90% of reptiles, even if healthy, can have Salmonella.

What are the symptoms of Salmonella?

Children are the most likely to get Salmonellosis. Most people infected with Salmonella develop diarrhoea, fever and abdominal cramps 12 to 72 hours after infection. The illness usually lasts for 4 to 7 days and most people recover without treatment.

However, in some people the diarrhoea may be severe enough to require admission to hospital.

Very severe infections can be fatal. The elderly, infants and those with impaired immune systems are more likely to have a severe illness.

Are there any longer term complications of Salmonellosis?

Most people recover quickly without treatment although it may be several months before bowel symptoms completely disappear.

Long term effects, which include a condition called Reiter's syndrome, are rare.

Some people continue to carry the germ although they themselves are well. In general, these people pass out the germ in small numbers and do not pose a risk to others.

Many different kinds of illnesses can cause diarrhoea, fever, or abdominal cramps. Salmonellosis can therefore only be diagnosed by growing Salmonella from a stool sample in a laboratory.

Once the Salmonella has been grown, further testing can determine its specific type and which antibiotics should be used to treat it, if the severity of the illness warrants it.

How is Salmonellosis treated?

Most Salmonellosis cases clear up without treatment. Because over-use of antibiotics breeds resistant germs, antibiotics are held back for severe cases.

Antibiotic treatment does not make a difference to whether the person develops longer term complications.

What can I do to prevent Salmonellosis? What can I do if I have it?

There is no vaccine to prevent Salmonellosis, nor do you become immune to it, so it is important to prevent it spreading and to avoid infection as far as possible.

- Do not eat raw or undercooked eggs, poultry or meat. You may not recognise that raw eggs could be present in home-made sauces, dressings, ice-creams, mousses and other desserts.
- Poultry and meat, including beef burgers, should be well cooked, not pink in the middle.
- If you are served undercooked poultry in a restaurant, do not eat it and complain to the manager and to your local Environmental Health Department.
- Do not consume unpasteurised milk.
- Wash produce thoroughly before consuming.
- Wash your hands with soap and hot water before and after handling raw meats.

Salmonella Infection Food Poisoning Fact Sheet 3

- Avoid cross-contamination of foods by keeping raw and cooked foods apart.
- Carefully clean all cutting boards, work surfaces and utensils with detergent and hot water after preparing raw meats. Use separate chopping boards for raw and ready-to-eat foods wherever possible.
- If you have the illness and work with children or the elderly, or work in a hospital or a food business, advise your supervisor. Your local Environmental Health Department, who can investigate to try and prevent further cases, may contact you.
- Salmonella germs stay in your system for at least several days after symptoms stop, so it is important to maintain a very high level of personal hygiene during this time.
- People who have Salmonellosis should not prepare food for others until they have been free of diarrhoea for at least 48hrs.

More Information about Salmonella

Salmonellas form a group of over 2000 bacteria that can cause diarrhoea in humans.

Salmonella enteritidis and typhimurium are the most common types of Salmonella to cause illness in the United Kingdom.

Salmonella has been known to cause illness for over 100 years. The germ was discovered by an American scientist called Salmon, after whom it was named.

- 1 Campylobacter
- 2 Rotavirus
- 3 Salmonella**
- 4 E. coli 0157
- 5 Clostridium perfringens
- 6 Shigella
- 7 Giardia
- 8 Viral Gastroenteritis



Council of the
ISLES OF SCILLY